

Toxicological assessment of inorganic arsenic and zinc content in button mushrooms

Soheil Sobhanardakani^{1,✉}, Amin Jahangard²

1. Department of the Environment, College of Basic Sciences, Hamedan Branch, Islamic Azad University, Hamedan, Iran
2. Young Researchers and Elite Club, Hamedan Branch, Islamic Azad University, Hamedan, Iran

Date of submission: 10 Jun 2017, **Date of acceptance:** 14 Jan 2018

ABSTRACT

Nowadays, discharge of toxic heavy metals into the environment is an increasing global concern. In light of this and that the per capita consumption of mushrooms has increased in Iran, the current study was carried out to investigate the levels and health-risk assessment of elements arsenic (As) and zinc (Zn) through consumption of button mushrooms marketed in Kermanshah city in 2016. In this descriptive study, a total of 30 samples from 10 brands of edible mushrooms were collected from the market basket of Kermanshah city. After preparation and processing of the samples in the laboratory, the contents of the studied elements were determined using ICP-OES. Also, all statistical analyses were performed using the SPSS version 19 statistical package. Based on the results, the contents (mg/kg) of arsenic and zinc with an average of 65.23 ± 13.57 and with an average of 66.23 ± 2.80 respectively were higher than the maximum permissible limit (MPL). Also, the results showed that all the computed values of health risk index (HRI) of both elements in adults, but only Zn in children, were within safe limits ($HRI < 1$). On the other hand, the HRI values of children with an average of 1.47 were more than the safe limit and led to a potential health risk under the current mushrooms consumption per capita for this age group. Therefore, serious attention to the discharge of chemicals into the environment and monitoring of the residue levels of pollutants, especially toxic heavy metals and pesticides in the foodstuffs, is recommended.

Keywords: mushroom; toxicological assessment; arsenic; zinc; health risk

Introduction

Since mushrooms are able to biodegrade the substrate and therefore use agricultural production wastes, they play an important role in the ecosystem. Fruiting bodies of mushrooms are valued for their texture, flavor, and nutritional properties. Also, uncontaminated mushrooms have been known as therapeutic foods, useful in preventing diseases including hypercholesterolemia, hypertension, and cancer.¹⁻³ Therefore, for these functional characteristics of mushroom, especially their ability to accumulate toxic heavy metals as common pollutants present in the environment, a survey of their chemical composition is

important.^{3,4}

Nowadays, to provide an adequate supply of food for the world's growing population and also in the control of pests and insect-borne diseases in crops, the use of chemical fertilizers, heavy metal-based pesticides, and organic fertilizers as an agricultural inputs is essential.⁵ Therefore, discharge of chemicals, particularly toxic heavy metals, into the environment in this way has caused an increasing global concern. Toxic heavy metals pollution is a major problem concerning the quality of the environment. Different metalloids or heavy metals such as As, Cd, Hg, Pb, and Ni are toxic, but, due to many elements such as Cr, Cu, Fe, Mn, and Zn known as enzyme activators, these groups of elements are vital for metabolic processes in low amounts. Nevertheless, these elements become toxic as the level increases.^{4,6}

Weathering of sedimentary and igneous rock is the major source of arsenic as a naturally-

✉ Soheil Sobhanardakani
s_sobhan@iauh.ac.ir

Citation: Sobhanardakani S, Jahangard A. Toxicological assessment of inorganic arsenic and zinc content in button mushrooms. J Adv Environ Health Res 2017; 5(4): 246-251

occurring element. However, the smelting of non-ferrous metals, mining activities, production of energy from fossil fuels, and also applications of agricultural inputs including arsenical pesticides or herbicides, are the anthropogenic origin of this element.⁷⁻⁹ Black foot disease, cancers of the kidney, liver, lung, bladder, prostate, colon, and skin are the most important diseases related to arsenic.^{10,11} However, zinc is known as an essential functional and structural element in several cellular processes, but an excessive intake of this element can be critical in its effects on growth, and plays a main role in cancer while also harming some physiological activities.^{12,13}

A review of the related literature has revealed the potential of mushrooms to accumulate pollutant compounds present in the environment at trace contents, particularly heavy metals.^{4,14-16} In this regard, the results of other studies have showed that the fruiting bodies of mushrooms have the ability to accumulate heavy metals.^{1-4,6,17-24}

The aim of risk assessment as part of risk analysis is a study of adverse health effects associated with exposure to a chemical. Therefore, identification and collection of information about the chemicals health hazards, exposure of humans to the hazardous substances, and the relationships between exposure, dose, and the serious impacts of chemicals on human health are the requisites of risk assessment.²⁵ Mushrooms form an important part of human diet in many countries in the world,⁴ and their consumption is growing rapidly in Iran with 880 g consumption per capita. Therefore, the aim of the current research is to investigate As and Zn concentrations of the fruiting bodies of button mushrooms collected in Kermanshah city, in order to provide information on the possible health risks associated with the consumption of mushrooms. This was done through evaluation of the daily intake of metal (DIM), the health risk index (HRI), and total HRI (THRI) of Ar and Zn, respectively.

Materials and Methods

Sample collection

In this study, 30 samples of 10 different brands of mushrooms (*Agaricus bisporus*) were purchased from different markets in

Kermanshah city in 2016 and were used for analysis of levels of As and Zn.

Chemical analyses

After washing the mushroom samples with distilled water, 1 g of the cap of each sample was placed in a porcelain crucible and ashed in an oven at 420–450 °C for 24 h. Then, ashed material was dissolved in 2 mL concentrated nitric acid (ultrapure grade, Sigma-Aldrich), evaporated to dryness, heated again for 3 h to 450 °C, dissolved in 3 mL mixture of concentrated nitric and sulfuric acid (2:1 v/v), and diluted with double distilled water (ddH₂O) up to 25 mL. Also, a blank digest was carried out in the same way. Finally, for the As and Zn analyses with three replications, a Varian710-ES inductively coupled plasma-optical emission spectrometry was used in this study.²⁶

Statistical analysis

The obtained results were statistically analyzed in the current study as follows: The normality of the distribution of data was analyzed using the Shapiro-Wilk test. Then the variance homogeneity was analyzed using an analysis of variance parametric test with a DMS post hoc and Duncan multiple range test. Also, the mean contents of elements were compared with the international standard using a one-sample test. The statistical calculations were done using statistical package SPSS version 20.0 (SPSS Inc., Chicago, IL, USA).

Potential health risk assessment

In this study, the risks to human health posed by chronic exposure to heavy metals were assessed. The average daily intake of metal was evaluated for the study of potential health risk assessment, in accordance with Eq. 1:²⁷

$$DIM = \frac{C_{\text{metal}} \times C_{\text{factor}} \times D_{\text{food intake}}}{\text{Baverage weight}} \quad (1)$$

where C_{metal} , C_{factor} , and $D_{\text{food intake}}$ represent the heavy metal concentrations in analyzed foodstuffs (mg/kg), conversion factor (0.085) and daily intake of mushrooms (2.40×10^{-3} kg per person per day), respectively.^{28,29} Also, $B_{\text{average weight}}$ indicates average body weight equaling 70.0 kg and 15.0 kg for adults and children, respectively.^{30,31}

The health risk index for button mushrooms' consumers was evaluated using Eq. 2:²⁷

$$HRI = \frac{DIM}{RfD} \quad (2)$$

In this equation, DIM indicates daily intake of metal (mg) and RfD refers to reference dose of metal (mg/kg/day). The oral reference doses were 0.0003 for As, and 0.30 for Zn. Also, an HRI<1 means the mushroom consumption is assumed to be safe.³²⁻³⁴

Total HRI (THRI) of elements for the mushrooms was evaluated in accordance with Eq. 3:²⁷

$$THRI = HRI (\text{toxicant 1}) + HRI (\text{toxicant 3}) + \dots + HRI (\text{toxicant n}) \quad (3)$$

Results and Discussion

Nowadays, the use of metal-containing pesticides, micronutrient and organic fertilizers, including arsenic-based pesticides, zinc sulfate and compost for the cultivation of mushrooms, can lead to the accumulation of trace elements and toxic heavy metals in their fruiting body. Based on the results, among the studied mushroom samples, arsenic was detected in amounts ranging from 40.0 mg/kg to 85.0 mg/kg, and zinc ranged from 61.0 mg/kg to 72.0 mg/kg (Table 1).

Table 1. Residual levels of examined elements in edible mushroom sample (mg/kg, dry weight)

Sample	Metal Concentration	
	As	Zn
1	74.33 ± 5.13 ^d	66.0 ± 4.36 ^a
2	63.33 ± 8.02 ^{bc}	67.33 ± 3.06 ^a
3	57.67 ± 5.86 ^b	66.33 ± 3.06 ^a
4	44.0 ± 5.29 ^a	64.33 ± 0.58 ^a
5	42.0 ± 2.64 ^a	68.0 ± 2.65 ^a
6	69.33 ± 3.79 ^{cd}	68.0 ± 5.29 ^a
7	77.33 ± 7.51 ^d	64.33 ± 0.58 ^a
8	77.67 ± 5.86 ^d	66.33 ± 1.53 ^a
9	75.0 ± 4.36 ^d	66.33 ± 0.58 ^a
10	71.67 ± 1.15 ^{cd}	65.33 ± 3.79 ^a
Min.	42.0	64.33
Max.	77.67	68.0
Mean	65.23	66.23
S.D.	13.57	2.80

Comparing the element contents in mushroom samples with the MPL (mg/kg) (0.10 for As and 50.0 for Zn) established by FAO/WHO^{35,36} showed that the mean contents of both elements observed in all samples were higher than the permissible limits. Duncan's test

results (different roman letters are presented in each column of Table 1) indicate statistical significant differences (*P*<0.05) in the contents of As and Zn between some brands of mushrooms. Based on the results, no statistically significant differences were observed in the Zn concentration between all brands (samples) of mushrooms.

In addition, all the calculated HRI values of arsenic and zinc in adults, and Zn in children were within the safe limits (HRI<1) in the current study (Table 2). Furthermore, the THRI values in edible mushrooms, which varied from 0.39 to 0.83 for adults and from 1.81 to 3.85 for children, were also within the safe limit (THRI<1) only for adults. Therefore, it can be concluded that, through only consuming button mushrooms under the mentioned consumption rate, children might have potential significant health risk.

Table 2. Daily intakes of metals (DIM, mg) for individual heavy metal

Brand	Adults		Children	
	As	Zn	As	Zn
1	0.0002	0.0002	0.001	0.0009
2	0.0002	0.0002	0.0009	0.0009
3	0.0002	0.0002	0.0008	0.0009
4	0.0001	0.0002	0.0006	0.0009
5	0.0001	0.0002	0.0006	0.0009
6	0.0002	0.0002	0.0009	0.0009
7	0.0002	0.0002	0.001	0.0009
8	0.0002	0.0002	0.001	0.0009
9	0.0002	0.0002	0.001	0.0009
10	0.0002	0.0002	0.001	0.0009

Table 3. Health risk index (HRI) for individual heavy metals caused by the button mushroom

Brand	Adults		Children	
	As	Zn	As	Zn
1	0.722	0.0006	3.37	0.003
2	0.615	0.0006	2.87	0.003
3	0.560	0.0006	2.61	0.003
4	0.427	0.0006	1.99	0.003
5	0.408	0.0007	1.90	0.003
6	0.673	0.0007	3.14	0.003
7	0.751	0.0006	3.51	0.003
8	0.754	0.0006	3.52	0.003
9	0.728	0.0006	3.40	0.003
10	0.696	0.0006	3.25	0.003

Since inorganic arsenic is acutely toxic, intake of large quantities of inorganic arsenic leads to some adverse health effects, including gastrointestinal symptoms, severe disturbances of the central nervous systems and cardiovascular disease, and eventually

death.^{26,37} In the present study, the mean contents of As in mushroom samples with 65.23 ± 13.57 mg/kg were much higher than the MPL. In this regard, Zhang et al. reported that the mean levels of As in caps of *Laccaria* mushrooms consumed in Yunnan, China were 87.80 ± 69.50 mg/kg.³⁸ Also, the results of other studies showed that the As contents (mg/kg) in caps of mushrooms consumed in Ljubljana, Slovenia, were in the range of 41–182,³⁹ in caps of mushrooms consumed in Domzale, Slovenia, they were in the range of 109–200,⁴⁰ and in caps of mushrooms consumed in Slovenia they were in the range of 0.80–129.⁴¹ Chen et al. also reported the content of As in the wild edible mushrooms from China being in the range of 0.44–1.48 mg/kg.²⁶

Zn has an important role in biological systems and plays an essential role in the structure of RNA, DNA, and protein synthesis, and also is necessary for the functioning of various enzymes that are known as essential nutrients. The major symptoms of deficiency of this element are slow maturation and delayed growth.²⁴ Based on the results of this study; the mean levels of Zn in caps of mushroom samples with 66.23 ± 2.80 mg/kg were generally lower than MPL. The mean contents of Zn (mg/kg) in the literature have been reported as 98.50 ± 34.75 in wild edible mushrooms in Soguksu National Park, Turkey,³ 38.83 ± 2.15 in *Pleurotus ostreatus*, 102.15 ± 33.28 in *Lycoperdon perlatum*, and 42.14 ± 2.45 in *Fistulina hepatica* from industrial areas of Romania,⁴ 4.35 ± 3.0 in wild mushrooms consumed in Zimbabwe,⁴² 64.62 ± 25.37 in wild edible mushroom samples consumed in Canakkale Province, Turkey,²⁴ and 43.84 ± 8.70 in wild edible mushrooms from Jordan.⁴³ Zinc content of mushroom samples in other study ranged from 30 mg/kg to 150 mg/kg.¹⁵

As shown in Table 2, the HRI values of arsenic and zinc for adults and the HRI values of Zn for children were less than 1, while the HRI value of As for children was higher than the safe limits. Here the average HRI value in edible mushroom samples was 0.316 for adults and 1.47 for children. Therefore, it can be admitted that children in the target population might have potential significant non-carcinogenic risks through only consuming analyzed button

mushrooms from Kermanshah city. In this regard, Nharingo et al. reported that the consumption by children of 300 g of fresh *Amanita loosii* and *Cantharellus floridulus* per day less than 16 kg harvested from Rail Block forest in Zimbabwe would cause health problems.⁴² Also, Zhang et al. after analyzing the As content in caps of *Laccaria* mushrooms consumed in China, reported that due to As levels being high in the fruit bodies of the studied samples, consumption of these foods may pose risks to human health.³⁸

Conclusions

Based on the results, the average HRI values of analyzed mushrooms for children of 1.47 were higher than the safe limits (HRI<1), and lead to potential health risks for children through consumption of mushrooms under the current consumption rate. Therefore, serious attention to the discharge of chemicals, especially toxic heavy metals and pesticides, to the environment and their content or residue monitoring in the foodstuffs, is recommended.

Acknowledgments

Authors are thankful to the Hamedan Branch, Islamic Azad University for providing instruments to conduct and complete this study.

References

1. Sesli E, Tuzen M, Soylak M. Evaluation of trace metal contents of some wild edible mushrooms from Black sea region, Turkey. *J Hazard Mater* 2008;160(2-3):462-7.
2. Ouzouni PK, Petridis D, Koller W-D, Kyriakos A, Riganakos KA. Nutritional value and metal content of wild edible mushrooms collected from West Macedonia and Epirus, Greece. *Food Chem* 2009;115(4):1575-80.
3. Sarikurkcü C, Copur M, Yildiz D, Akata I. Metal concentration of wild edible mushrooms in Soguksu National Park in Turkey. *Food Chem* 2011;128(3):731-4.
4. Radulescu C, Stihi C, Busuioc G, Gheboianu AI, Popescu IV. Studies concerning heavy metals bioaccumulation of wild edible mushrooms from industrial area by using spectrometric techniques. *Bull Environ Contam Toxicol* 2010;84(5):641-6.
5. Sobhanardakani S. Residual levels of diazinon and benomyl on greenhouse mushrooms. *Iran J Toxicol* 2015;9(29):1307-11.
6. Yamaça M, Yıldız D, Sarikürkcü C, Çelikkollu M, Halil Solak M. Heavy metals in some edible

- mushrooms from the Central Anatolia, Turkey. *Food Chem* 2007;103(2):263-7.
7. Stolz JF, Basu P, Santini JM, Oremland RS. Arsenic and selenium in microbial metabolism. *Annu Rev Microbiol* 2006;60:107-30.
 8. Abu Bakar AF, Yusoff I, Fatt NT, Ashraf MA. Cumulative impacts of dissolved ionic metals on the chemical characteristics of river water affected by alkaline mine drainage from the Kuala Lipis gold mine, Pahang, Malaysia. *J Chem Ecol* 2015;31(1):22-33.
 9. Alizamir M, Sobhanardakani S. Forecasting of heavy metals concentration in groundwater resources of Asadabad Plain using artificial neural network approach. *J Adv Environ Health Res* 2016;4(2):68-77.
 10. Tseng CH, Tai TY, Chong CK, Tseng CP, Lai MS, Lin BJ, et al. Long-term arsenic exposure and incidence of non-insulin-dependent diabetes mellitus: A cohort study in arseniasis-hyperendemic villages Taiwan. *Environ Health Perspect* 2000;108(9):847-51.
 11. Liang CP, Wang SW, Kao YH, Chen JS. Health risk assessment of groundwater arsenic pollution in southern Taiwan. *Environ Geochem Health* 2016;38(6):1271-81.
 12. Grattan BJ, Freake HC. Zinc and cancer: Implications for LIV-1 in breast cancer. *Nutrients* 2012;4(7):648-75.
 13. Medeiros RJ, Dos Santos LMG, Freire AS, Santelli RE, Braga AMCB, Krauss TM, et al. Determination of inorganic trace elements in edible marine fish from Rio de Janeiro State, Brazil. *Food Control* 2012;23(2):535-41.
 14. Kalač P, Svoboda L, Havlíčková B. Contents of cadmium and mercury in edible mushrooms. *J Appl Biomed* 2004;2:15-20.
 15. Kalač P, Svoboda L. A review of trace element concentrations in edible mushrooms. *Food Chem* 2000;69 (3):273-81.
 16. Ita BN, Essien JP, Ebong GA. Heavy metal levels in fruiting bodies of edible and non-edible mushrooms from the Niger Delta Region of Nigeria. *J Agric Soc Sci* 2006;2(2):84-7.
 17. Svoboda L, Kalac P. Contamination of two edible *Agaricus* spp. mushrooms growing in a town with cadmium, lead, and mercury. *Bull Environ Contam Toxicol* 2003;71(1):123-30.
 18. Turkekul I, Elmastas M, Tuzen M. Determination of iron, copper, manganese, zinc, lead, and cadmium in mushroom samples from Tokat, Turkey. *Food Chem* 2004;84(3):389-92.
 19. Soylak M, Saracoglu S, Tuzen M, Mendil D. Determination of trace metals in mushroom samples from Kayseri, Turkey. *Food Chem* 2005;92(4):649-52.
 20. Svoboda L, Havlickova B, Kalac P. Contents of cadmium, mercury and lead in edible mushrooms growing in a historical silver-mining area. *Food Chem* 2006; 96(4): 580-5.
 21. Tuzen M, Sesli E, Soylak M. Trace element levels of mushroom species from East Black Sea region of Turkey. *Food Control* 2007; 18(7): 806-10.
 22. Antonijevic MM, Maric M. Determination of the content of heavy metals in pyrite contaminated soil and plants. *Sensors (Basels)* 2008;8(9):5857-65.
 23. Gencelep H, Uzun Y, Tuncturk Y, Demirel K. Determination of mineral contents of wild-grown edible mushrooms. *Food Chem* 2009;113(4):1033-6.
 24. Çayır A, Coşkun M, Coşkun M. The heavy metal content of wild edible mushroom samples collected in Canakkale Province, Turkey. *Biol Trace Elem Res* 2010;134(2):212-9.
 25. Sobhanardakani S. Health risk assessment of As and Zn in canola and soybean oils consumed in Kermanshah, Iran. *J Adv Environ Health Res* 2016;4(2):62-7.
 26. Chen XH, Zhou HB, Qiu GZ. Analysis of several heavy metals in wild edible mushrooms from regions of China. *Bull Environ Contam Toxicol* 2009; 83(2):280-85.
 27. Guo J, Yue T, Li X, Yuan Y. Heavy metal levels in kiwifruit orchard soils and trees and its potential health risk assessment in Shaanxi, China. *Environ Sci Pollut Res Int* 2016;23(14):14560-6.
 28. Sobhanardakani S, Sadri S, Jameh Bozorgi S. Evaluation of organophosphorus pesticide diazinon residue in greenhouse crops using spectrophotometry (case study: mushroom). *Food Hyg* 2004;3(4):73-80.
 29. Abdi S, Sobhanardakani S, Jameh Bozorgi S. Monitoring of benomyl residue in mushroom marketed in Hamadan City. *Sci J Hamadan Univ Med Sci* 2015; 22(2): 137-43.
 30. Omar WA, Zaghoul KH, Abdel-Khalek AA, Abo-Hegab S. Risk assessment and toxic effects of metal pollution in two cultured and wild fish species from highly degraded aquatic habitats. *Arch Environ Contam Toxicol* 2013;65(4):753-64.
 31. Tang W, Cheng J, Zhao W, Wang W. Mercury levels and estimated total daily intakes for children and adults from an electronic waste recycling area in Taizhou, China: Key role of rice and fish consumption. *J Environ Sci (China)* 2015;34:107-15.
 32. Xue ZJ, Liu SQ, Liu YL, Yan YL. Health risk assessment of heavy metals for edible parts of vegetables grown in sewage-irrigated soils in suburbs of Baoding City, China. *Environ Monit Assess* 2012;184(6):3503-13.

33. Liang Q, Xue ZJ, Wang F, Sun ZM, Yang ZX, Liu SQ. Contamination and health risks from heavy metals in cultivated soil in Zhangjiakou City of Hebei Province, China. *Environ Monit Assess* 2015;187(12):754.
34. Zeng X, Wang Z, Wang J, Guo J, Chen X, Zhuang J. Health risk assessment of heavy metals via dietary intake of wheat grown in Tianjin sewage irrigation area. *Ecotoxicology* 2015;24(10):2115-24.
35. Taghipour H, Mosafieri M. Heavy metals in the vegetables collected from production sites. *Health Promot Perspect* 2013;3(2):185-93.
36. Ogunkunle ATJ, Bello OS, Ojofeitimi OS. Determination of heavy metal contamination of street-vended fruits and vegetables in Lagos state, Nigeria. *Int Food Res J* 2014;21(5):1725-30.
37. Järup L. Hazards of heavy metal contamination. *Br Med Bull* 2003;68:167-82.
38. Zhang J, Li T, Yang Y-L, Liu H-G, Wang Y-Z. Arsenic concentrations and associated health risks in *Laccaria* mushrooms from Yunnan (SW China). *Biol Trace Elem Res* 2015;164(2):261-6.
39. Byrne AR, Tušek-Žnidarič M. Arsenic accumulation in the mushroom *Laccaria amethystina*. *Chemosphere* 1983;12(7-8):1113-17.
40. Byrne AR, Tušek-Žnidarič M, Puri BK, Irgolic KJ. Studies of the uptake and binding of trace metals in fungi Part II Arsenic compounds in *Laccaria amethystina*. *Appl Organomet Chem* 1991; 5(1):25-32.
41. Slekovec M, Irgolic KJ. Uptake of arsenic by mushrooms from soil. *Chem Spec Bioavailab* 1996;8(3-4):67-73.
42. Nharingo T, Ndumo T, Moyo M. Human health risks due to heavy metals through consumption of wild mushrooms from Macheke forest, Rail Block forest and Muganyi communal lands in Zimbabwe. *Environ Monit Assess* 2015;187(12):738.
43. Semreen MH, Aboul-Enein HY. Determination of heavy metal content in wild-edible mushroom from Jordan. *Anal Lett* 2011;44(5):932-41.